

BARNSTEAD SCHOOL DISTRICT
Universal Precautions & Infection Control

Recent concern about how children with AIDS should be educated has raised several questions regarding exposure of teachers and children to potentially infectious body fluids from children with communicable diseases in the school setting:

1. Does contact with body fluids present a risk of infection?
2. What should be done to avoid contact with potentially infected body fluids?
3. What should be done if direct contact with body fluids is made?
4. How should such fluids when spilled be removed from the environment?

The following guidelines are meant to provide simple and effective precautions against transmission of disease for all persons, including pregnant women, potentially exposed to the blood or body fluids of any student. No distinction is made between body fluids from students with a known disease or those from students without symptoms or with an undiagnosed disease.

DOES CONTACT WITH BODY FLUIDS PRESENT A RISK?

The body fluids of all persons should be considered to contain potentially infectious agents (germs). The term "body fluids" includes: blood, semen, drainage from scrapes and cuts, feces, urine, vomitus, respiratory secretions (e.g., nasal discharge) and saliva. Contact with body fluids presents a risk of infection with a variety of germs. In general, however, the risk is very low and dependent on a variety of factors including the type of fluid with which contact is made and the type of contact made with it.

It must be emphasized that with the exception of blood, which is normally sterile, the body fluids with which one may come in contact usually contain many organisms, some of which may cause disease. Furthermore, many germs may be carried by individuals who have no symptoms of illness. These individuals may be at various stages of infection: incubating disease, mildly infected without symptoms, or chronic carriers of certain infectious agents including the AIDS and hepatitis viruses. In fact, transmission of communicable diseases is more likely to occur from contact with fluids from unrecognized individuals because simple precautions are not always carried out.

WHAT SHOULD BE DONE TO AVOID CONTACT WITH BODY FLUIDS?

When possible, direct skin contact with body fluids should be avoided. Disposable gloves are recommended when direct hand contact with body fluids is anticipated (e.g., treating bloody noses, handling clothes soiled by incontinence, cleaning small spills by hand). Gloves used for this purpose should be put in a plastic bag or lined trash can, secured, and disposed of daily. Hands should be washed thoroughly after any contact with body fluids and before contact with any other person.

WHAT SHOULD BE DONE IF DIRECT SKIN CONTACT OCCURS?

In many instances, unanticipated skin contact with body fluids may occur in situations where gloves may be immediately unavailable (e.g., when wiping a runny nose, applying pressure to a bleeding injury outside the classroom, helping a child in the bathroom). In these instances, hands and other affected skin areas of all exposed persons should be routinely washed with soap and water after direct contact has ceased. Clothing and other nondisposable items (e.g., towels used to wipe up body fluid) that are soaked through with body fluids should be rinsed and placed in plastic bags. If presoaking is required to remove stains, (e.g., blood, feces), use gloves to rinse or soak the item in cold water prior to bagging. Clothing should be sent home for washing with appropriate directions to parents/teachers (see page 6). Disposable towels and tissues should be used whenever possible. Contaminated disposable items (e.g., tissues, paper towels, diapers) should be handled as with disposable gloves.

HOW SHOULD SPILLED BODY FLUIDS BE REMOVED FROM THE ENVIRONMENT?

Most schools have standard procedures already in place for removing body fluids (e.g., vomitus). These procedures should be reviewed to determine whether appropriate cleaning and disinfection steps have been included. Many schools stock sanitary absorbent agents specifically intended for cleaning body fluid spills (e.g., ZGOOP, Parsen Mfg. Co., Philadelphia, PA). Disposable gloves should be worn when using these agents. The dry material is applied to the area, left for a few minutes to absorb the fluid, and then vacuumed or swept up. The vacuum bag or sweepings should be disposed of in a plastic bag. Broom and dustpan should be rinsed in a disinfectant. No special handling is required for vacuuming equipment.

HANDWASHING PROCEDURES

Proper handwashing requires the use of soap and water and vigorous washing under a stream of running water for approximately 10 seconds. Soap suspends easily removable soil and microorganisms allowing them to be washed off. Running water is necessary to carry away dirt and debris. Rinse under running water. Use paper towels to thoroughly dry hands.

DISINFECTANTS

An intermediate level of disinfectant should be used to clean surfaces contaminated with body fluids. Such disinfectants will kill vegetative bacteria, fungi, tubercle bacillus and viruses. The disinfectant should be registered by the U.S. Environmental Protection Agency (EPA) for use as a disinfectant in medical facilities and hospitals.

Various classes of disinfectants are listed below. Hypochlorite solution (bleach) is preferred for objects that may be put in the mouth.

1. Ethyl or isopropyl alcohol (70%)
2. Phenolic germicidal detergent in a 1% aqueous solution (e.g., Lysol*).

3. Sodium Hypochlorite with at least 100 ppm available chlorine (1/2 cup household bleach in 1 gallon water, needs to be freshly prepared each time it is used).
4. Quarternary ammonium germicidal detergent in 2% aqueous solution (e.g., Triquat*, Mytar* or Sage*).
5. Iodophor germicidal detergent with 500 ppm available iodine (e.g., Wescodyne*).

DISINFECTION OF HARD SURFACES AND CARE OF EQUIPMENT

After removing the soil, a disinfectant is applied. Mops should be soaked in the disinfectant after use and rinsed thoroughly or washed in a hot water cycle before rinse. Disposable cleaning equipment and water should be placed in a toilet or plastic bag as appropriate. Non-disposable cleaning equipment (dust pans, buckets) should be thoroughly rinsed in the disinfectant. The disinfectant solution should be promptly disposed down a drain pipe. Remove gloves and discard in appropriate receptacles.

DISINFECTION OF RUGS

Apply sanitary absorbent agent, let dry and vacuum. If necessary, mechanically remove with dust pan and broom, then apply rug shampoo (a germicidal detergent) with a brush and revacuum. Rinse dust pan and broom in disinfectant. If necessary, wash brush with soap and water. Dispose of nonreusable cleaning equipment as noted above.

LAUNDRY INSTRUCTIONS FOR CLOTHING SOILED WITH BODY FLUIDS

The most important factor in laundering clothing contaminated in the school setting is elimination of potentially infectious agents by soap and water. Addition of bleach will further reduce the number of potentially infectious agents. Clothing soaked with body fluids should be washed separately from other items. Presoaking may be required for heavily soiled clothing. Otherwise, wash and dry as usual. If the material is bleachable, add 1/2 cup household bleach to the wash cycle. If material is not colorfast add 1/2 cup nonchlorox bleach (e.g., Clorox II, Borateem) to the wash cycle.

*Brand names used only for examples of each type of germicidal solution and should not be considered an endorsement of a specific product.